arXiv:gr-qc/0007084v2 25 Oct 2000

IMSc/2000/07/37

Quasi-normal modes of AdS black holes : A superpotential approach

T.R.Govindarajan [1] and V.Suneeta [2]

The Institute of Mathematical Sciences, CIT Campus, Chennai 600113, India

Abstract

A novel method, based on superpotentials is proposed for obtaining the quasi-normal modes of anti-deSitter black holes. This is inspired by the case of the three dimensional BTZ black hole, where the quasi-normal modes can be obtained exactly and are proportional to the surface gravity. Using this approach, the quasi-normal modes of the five dimensional Schwarzschild anti-deSitter black hole are computed numerically. The modes again seem to be proportional to the surface gravity for very small and very large black holes. They reflect the well-known instability of small black holes in anti-deSitter space.

PACS No.: 04.62.+v, 04.70.-s

---

[1] e-mail: trg@imsc.ernet.in

[2] e-mail: suneeta@imsc.ernet.in

1 Introduction

Numerical studies of perturbations of black holes have revealed the presence of certain characteristic modes that govern the time evolution of the initial perturbation. These modes seem to depend only on the black hole parameters and not on the nature of the perturbation. This was first recognised by Vishveshvara [1] while studying perturbations of Schwarzschild black holes. Since then, perturbations of asymptotically flat black holes have been analysed and these characteristic modes(called quasi-normal modes) have been found for some black holes (a detailed account can be found in [2], [3]).

Recently, there has been an interest in quasi-normal modes of AdS black holes in light of the AdS/CFT correspondence [4]. There is some evidence to suggest that an off-equilibrium configuration in the bulk AdS space is related to an off equilirium state in the boundary theory. In [5], black hole formation by collapse of a thin shell in AdS space is investigated. The shell is related by AdS/CFT duality to an off-equilibrium state in the boundary theory which evolves towards equilibrium when the shell collapses to form a black hole.

A quasi-normal mode governs the decay in time of a perturbation of the black hole configuration in the bulk. It should therefore be related by AdS/CFT duality to the return of the boundary Yang-Mills theory to thermal equilibrium. The role of ingoing modes in determining the thermalization time scale of the boundary Yang-Mills theory was pointed out in [6][3]. A numerical computation of quasi-normal mode frequencies for AdS black holes in various dimensions has been done in [7].

In this paper, we compute the quasi-normal modes for the AdS-Schwarzschild black hole in five dimensions using a superpotential approach. This is done for the perturbation of a black hole by a minimally coupled scalar field. Time-independent normal modes of a scalar field in the background of this black hole have already been analysed [8, 9].

We notice, that for the $(2+1)-d$ BTZ black hole, exact quasi-normal mode solutions can be obtained from the Klein-Gordon equation as the black hole potential belongs to a class of exactly solvable potentials, derived from a superpotential.

We propose, based on analogy with the results for the $(2+1)-d$ BTZ black hole, and on numerical evidence in five dimensions, that the black hole potential can be described by a potential series derived from a superpotential. The problem is then exactly solvable and the form of the quasi-normal mode wave function is thus known. This is used as an ansatz to obtain the quasi-normal modes of the AdS_5 black hole. However, the numerical values of the mode frequencies do not agree with the earlier calculation in [7]. We comment on this discrepancy.

---

[3] The normalisation measure used here is not appropriate, however, and these modes are not quasi-normal modes.

The organisation of this paper is as follows: The next section is a brief review of quasi-normal modes and their properties, and a short summary of some numerical approaches to computing these modes.

The actual computation of quasi-normal(QN) modes for the AdS_5-Schwarzschild black hole is described in Section 3. First, the QN modes for the $(2+1)-d$ BTZ black hole are evaluated. Motivated by this, a superpotential approach to finding the QN modes for the five-dimensional black hole is presented and the mode frequencies obtained.

In Section 4, we discuss our results. We show that the behaviour of the modes computed numerically by us for small black holes is consistent with that expected from the differential equation obeyed by the mode solutions. We mention salient points of two earlier papers [6] and [7] on the subject and comment on the discrepancy between the numerical values obtained by us and those in [7]. We also state work in progress on QN modes for the RN AdS_5 black hole, and black holes in other dimensions.

2 Quasi-normal modes and their properties

Quasi-normal modes of a black hole are characteristic modes associated with the decay of any perturbation outside a black hole. In general, these modes do not form a complete set, in the sense that the time decay of a perturbation cannot be described *completely* in terms of them. However, they dominate the decay at certain intermediate or late times. To compute these modes, one studies the decay of an initial perturbation by imposing ingoing boundary conditions at the horizon and (for asymptotically flat black holes) outgoing boundary conditions at infinity. This ensures that no gravitational wave from the horizon or infinity disturbs the initial perturbation. Due to these boundary conditions, the quasi-normal modes are complex. The mode wave functions are non-normalisable in space but exponentially decay in time. Since the exact mode wave functions are not known even for the Schwarzschild black hole, there exist only numerical computations of quasi-normal modes. These are also very difficult to do as it is not easy to isolate the purely ingoing wave near the horizon that is also outgoing at infinity. Such a wave function blows up at these points and could be contaminated by a small outgoing part at the horizon or a small ingoing part at infinity where these parts go exponentially to zero. A unique boundary condition to isolate quasi-normal modes has been given by Nollert [2]. The QN modes are poles of the Green's function corresponding to the wave equation obeyed by the perturbing field. Using this, the QN modes for the Schwarzschild and Reissner-Nordstrom black holes were evaluated numerically [10], [11]. WKB methods have also been used to calculate these modes [12] (see also [2] for more references).

The method of continued fractions [13] gives very accurate results for the modes. Here, after removing the singular parts at the horizon and infinity, the QN mode wave function is expanded as a series. The coefficients in the series obey a three-term recurrence relation and depend on the quasi-normal mode frequency. The frequencies then have to satisfy a continued fraction relation. An interesting analytic approach to computing the QN modes is by approximating the black hole potential by an exactly solvable potential (the Poschl Teller or Eckart potential) [14], [15]. Ferrari and Mashhoon [14] have mapped the problem of QNMs of the Poschl-Teller potential to the bound states of the inverse potential and computed the modes. These approximate the fundamental QN modes of the Schwarzschild black hole well.

For the case of asymptotically anti-deSitter black holes, the black hole potential diverges at infinity. Therefore, as pointed out in [7], the quasi-normal modes are defined to be those that are ingoing at the horizon, but fall off to zero at infinity. It can be shown that this also agrees with the unique boundary condition of Nollert [2] for obtaining quasi-normal modes. However, due to the divergence of the AdS black hole potential at infinity, most of the methods described above to compute the QN modes of asymptotically flat black holes cannot be used directly. It is seen that a combination of two methods, inspired by the case of the $(2+1)-d$ black hole, yields good results.

3 Numerical computation of quasi-normal modes

We first consider the case of the QN modes of the non-rotating $(2+1)-d$ BTZ black hole [16]. The non-rotating BTZ black hole metric for a spacetime with negative cosmological constant $\Lambda = -\frac{1}{l^2}$ is given by

$$ds^2 \quad = \quad -(N)^2\ dt^2 + (N)^{-2}\ dr^2 + r^2\ (d\phi)^2 \tag{1}$$

with

$$N \;=\; \sqrt{(-M \;+\; \frac{r^2}{l^2})} \tag{2}$$

The horizon radius $r_+ \;=\; \sqrt{M}l$. We consider a massless scalar field in the black hole background. The Klein-Gordon equation for the scalar field is written using an ansatz (independent of the angular variable) for the field

$$\Phi = \frac{1}{\sqrt{r}}\ \chi(r)\ \ \exp(i\omega t) \tag{3}$$

and by going to the tortoise coordinate r_* where $dr_* = \frac{dr}{N^2}$. The Klein- Gordon equation is

$$-\frac{d^2\chi}{dr_*^2} \;+\; V(r)\chi \;=\; \omega^2\chi \tag{4}$$

where

$$V \;=\; \frac{3r^2}{4l^4} \;-\; \frac{M}{2l^2} \;-\; \frac{M^2}{4r^2} \tag{5}$$

Since $r = -r_+ \coth(\alpha r_*)$ where

$$\alpha \;=\; \frac{1}{2}\,\frac{d(N^2)}{dr}(r_+) \;=\; \frac{\sqrt{M}}{l} \tag{6}$$

$$V \;=\; \frac{M}{4l^2}\,(\frac{3}{(\sinh(\alpha r_*))^2} \;+\; \frac{1}{(\cosh(\alpha r_*))^2}) \tag{7}$$

α is the surface gravity of the black hole and is equal to $2\pi T$ where T is the Hawking temperature.

This potential can be obtained from a superpotential [17] of the form

$$W \;=\; A\,\coth(\alpha r_*) \;+\; B\,\tanh(\alpha r_*) \tag{8}$$

by

$$V \;=\; W^2 \;-\; (W)' \;-\; A^2 \;-\; B^2 \;-\; 2AB \tag{9}$$

Here, from (7),

$$A \;=\; -\,\alpha(1/2 \pm 1) \tag{10}$$

$$B \;=\; -\,\frac{\alpha}{2} \tag{11}$$

As discussed in [17], the lowest energy state for the potential $W^2 - (W)'$ has energy zero, so the lowest energy for the potential V is $-(A+B)^2$. The wave function corresponding to this state is

$$\chi \;=\; \exp(-\int(W)) \;\;=\;\; (\sinh(\alpha r_*))^{-A/(\alpha)}\;\;(\cosh(\alpha r_*))^{-B/(\alpha)} \tag{12}$$

For any of the two values of A and B in (11), the wave function (12) is not normalisable. It blows up either at the horizon or at $r = \infty$ (i.e $r_* = 0$). We are interested in the quasi-normal mode solution that blows up at the horizon and falls off to zero at infinity. This corresponds to $A = -(3/2)\alpha$. Also, $B = -\frac{\alpha}{2}$. The lowest energy (formally, since the wave function corresponding to this is non-normalisable) is then

$$E \;=\; -\,4(\alpha)^2 \tag{13}$$

and the lowest quasi-normal mode $\omega = \sqrt{E}$. Therefore,

$$\omega \quad = \quad -\,2\,i\,\alpha \tag{14}$$

Thus, for a given horizon radius $r_+ = \sqrt{M}l$, we can calculate α taking $l = 1$. Then we can read off the value of the lowest quasi-normal mode.

The case of the quasi-normal modes of the AdS_5-Schwarzschild black hole is not so simple as the black hole potential is more complicated than the BTZ case.

The metric for this black hole is

$$ds^2 \quad = \quad -\,(N)^2\;dt^2 + (N)^{-2}\;dr^2 + r^2\;(d\Omega_3)^2 \tag{15}$$

with

$$N \quad = \quad \sqrt{1\;+\;(r^2)/(l^2)\;-\;(r_0/r)^2} \tag{16}$$

where the black hole mass M is

$$M = \frac{3A_3 r_0^2}{16\pi G_5} \tag{17}$$

and A_3 is the area of a unit 3-sphere. The Klein-Gordon equation for the scalar field in the background of this black hole can be written, as before, as a potential problem. Using the ansatz for the field

$$\Phi \quad = \quad \left(\frac{1}{r}\right)^{3/2}\;\chi(r)\;\;\exp(i\omega t) \tag{18}$$

and by changing to the tortoise coordinate given by $dr_* = \frac{dr}{N^2}$, the Klein Gordon equation is

$$-\,\frac{d^2\chi}{dr_*^2}\quad +\quad V(r)\,\chi \quad = \quad \omega^2\,\chi \tag{19}$$

where

$$V(r) \quad = \quad \left(\frac{15}{4} + \frac{3}{4r^2} + \frac{9r_0^2}{4r^4}\right)\left(1 + r^2 - (\frac{r_0}{r})^2\right) \tag{20}$$

Here, for simplicity, we have taken $l = 1$. We have also not considered an angular dependence for the field, which is a simple generalisation of the following analysis.

If the potential could be written as an exactly solvable potential in the r_* coordinate, then the quasi-normal modes could be obtained easily from the 'lowest' energy, as in the BTZ case. As mentioned in the previous section, the fundamental QN modes of the Schwarzschild black hole were closely approximated by the QN modes of the Poschl-Teller potential. This is also an exactly solvable potential derivable from a superpotential [17].

We propose that the potential can be written as a series, derived from a superpotential of the form

$$W \;=\; \sum_{n=1}^{\infty} A_n \coth(n\alpha r_*) \tag{21}$$

where as in (6), α is the black hole surface gravity,

$$\alpha \;=\; (1/2)\,\frac{d(N^2)}{dr}(r_+) \;=\; r_+ \;+\; \frac{r_0^2}{r_+^3} \tag{22}$$

As before, $\alpha = 2\pi T$, where T is the Hawking temperature.

The potential derived from this series as in (9) is

$$V_s \;=\; \sum_{n=1}^{\infty} \frac{(A_n^2 + nA_n\alpha)}{\sinh^2(n\alpha r_*)} \;+\; \sum_{n,m=1,\; n\neq m}^{\infty} \frac{A_n A_m}{\tanh(n\alpha r_*)\tanh(m\alpha r_*)} \tag{23}$$

It has lowest energy

$$E \;=\; -\,(\sum_{n=1}^{\infty} A_n)^2 \tag{24}$$

The form of the potential is such that it goes exponentially to zero in r_* as $r \to r_+$ and blows up as $1/r_*^2$ as $r_* \to 0$ (or $r \to \infty$). This reproduces the behaviour of the black hole potential. We propose that this potential series is exactly equal to the black hole potential (20). Terminating this potential series at finite order would then mean an approximation to the black hole potential in the spirit of the Poschl-Teller method for asymptotically flat black holes. The wave function corresponding to the lowest energy state is

$$\chi \;=\; \exp(-\int(W)) \;\;=\;\; \prod_{n=1}^{\infty} (\sinh(n\alpha r_*))^{-A_n/(n\alpha)} \tag{25}$$

It is non-normalisable, as in the $(2+1)-d$ case. In order to calculate the quasi-normal modes for the AdS-Schwarzschild black hole, we use (25) as an ansatz in the equation (19). However, we truncate the product in the ansatz upto some finite order. This is equivalent to truncating the potential series. We first write

$$\chi \;=\; \psi\;\exp(-i\omega r_*) \tag{26}$$

This isolates the non-normalisable part of the wave function. We can rewrite (19) as an equation for ψ as

$$- \frac{d^2\psi}{dr_*^2} \; + \; 2i\omega\frac{d\psi}{dr_*} \; + \; V(r)\psi \; = \; 0 \tag{27}$$

Here, V is the black hole potential (20). We now try to solve for ω by taking the ansatz

$$\psi \; = \; \prod_{n=1}^{N}(1 \; - \; \exp(2n\alpha r_*))^{-\frac{A_n}{n\alpha}} \tag{28}$$

This ansatz is obtained from (25) which was an ansatz for χ. Thus, N represents the order upto which V_s has been truncated.

The ansatz for ψ is substituted into (27) and the l.h.s of (27) is expanded as a series in $(x - x_+)$ where

$$\begin{aligned} x \; &= \; \frac{1}{r} \\ x_+ \; &= \; \frac{1}{r_+} \end{aligned} \tag{29}$$

From (27), each term of the series is to be equated to zero. We first divide (27) by ψ. Then, we have

$$\begin{aligned} &- \; 4(\sum_{m=1}^{N} A_m K_m)^2 \; - \; 4 \; \alpha \; (\sum_{m=1}^{N} m A_m K_m) \; - \\ &4 \; \alpha \; (\sum_{m=1}^{N} m A_m K_m^2) \; + \; 4 \; i\omega \; (\sum_{m=1}^{N} A_m K_m) \; + \; V \; = \; 0 \end{aligned} \tag{30}$$

where

$$K_m \; = \; \frac{\exp(2m\alpha r_*)}{1 \; - \; \exp((2m\alpha r_*)} \tag{31}$$

Expanding each of the terms in the l.h.s of (30),

$$K_m \; = \; \sum_{n=0} c_m(n) \; (x - x_+)^n \tag{32}$$

$$K_m^2 \; = \; \sum_{n=0} d_m(n) \; (x - x_+)^n \tag{33}$$

$$V \; = \; \sum_{n=0} V_n \; (x - x_+)^n \tag{34}$$

Substituting these expansions into (30), and equating the term of order $(x-x_+)$ in the l.h.s to zero, we obtain

$$A_1 \;=\; -\;\frac{\alpha(\frac{15}{4} \;+\; \frac{3x_+^2}{4} \;+\; \frac{9r_0^2x_+^4}{4})}{2(i\omega \;-\; \alpha)c_1(1)} \tag{35}$$

Similarly, equating the term of order $(x-x_+)^N$ in the l.h.s to zero, we obtain a recursion relation for A_N.

$$\begin{aligned} A_N \;=\; & \frac{-\alpha(\sum_{m=1}^{N-1} mA_mc_m(N))}{(N\alpha \;-\; i\omega)c_N(N)} \;-\; \frac{\alpha(\sum_{m=1}^{N-1} mA_md_m(N))}{(N\alpha \;-\; i\omega)c_N(N)} \\ & -\; \frac{(\sum_{m,n=1}^{N-1} \;\; \sum_{N1,N2;N1+N2=N} A_mA_nc_m(N1)c_n(N2))}{(N\alpha \;-\; i\omega)c_N(N)} \\ & +\; i\omega\;\frac{(\sum_{m=1}^{N-1} A_mc_m(N))}{(N\alpha \;-\; i\omega)c_N(N)} \;+\; \frac{V_N}{4(N\alpha \;-\; i\omega)c_N(N)} \end{aligned} \tag{36}$$

The A_n coefficients are functions of ω and of the black hole parameters. The recursion relation for the A_n is complicated, unlike the case of the method of partial fractions [13] applied to asymptotically flat black holes. There, an ansatz was taken for the wave function (removing its singular part) and it was expanded as a series whose coefficients obeyed a simple three-term recursion relation. From (27), as $r_* \rightarrow 0$ (i.e as $r \rightarrow \infty$), the two kinds of solutions are $\psi \sim r_*^{\frac{-3}{2}}$ and $\psi \sim r_*^{\frac{5}{2}}$. The first solution is not normalisable. We want the quasi-normal mode solution to vanish as $r_* \rightarrow 0$. So we choose the second solution. But from our ansatz (28) for the solution, this implies a relation between the coefficients A_n. More precisely,

$$\sum_{n=1}^{N} \frac{A_n}{n\alpha} \;=\; -\;5/2 \tag{37}$$

Since the A_n coefficients are functions of ω, this relation gives the value of the lowest quasi-normal mode as a function of the black hole parameters. This would then be an approximation to the actual QN mode of the black hole at order N. Thus, the asymptotic behaviour of the actual solution is *explicitly* demanded of the ansatz. This is *used* to obtain the mode frequencies from the relation (37).

We have used this method to calculate the QN modes at various orders N. We find that as we increase N, there is a convergence in the mode frequency. The mode frequencies are given in Table 1. They have been calculated at order $N = 18$. Beyond this order, as (37) is

Radius r_+	Re(ω)	Im(ω)
1	0.6948	1.4648
2	1.0713	1.9817
5	2.4462	4.2642
10	4.8249	8.3279
50	24.0159	41.3183
100	48.0251	82.6165
150	72.0358	123.9190
500	240.1150	413.0500
750	360.1720	619.5740
1000	480.2290	826.0980

Table 1.

complicated, it becomes computationally time-consuming to find its roots. However, as there is a clear convergence in the mode frequency as one increases the order from $N = 1$ and as the percentage difference between the real and imaginary parts of the mode frequency at order $N = 18$ and $N = 17$ is about 1% (for $Re\omega$, it is 1.26%, and for $Im\omega$, it is 1.09%), these numbers should be a good approximation to the QN modes of the black hole (more discussion on results is given in the next section).

4 Discussion of results

The lowest quasi-normal mode frequencies for different horizon radii r_+ are given in Table 1. As mentioned in the last section, there is a convergence in the mode frequency as the order N is increased. The convergence curve for $r_+ = 10$ is given in Fig.1. The rate of convergence does not seem to depend on the value of the horizon radius r_+.

The real and imaginary parts of the mode frequencies are plotted as a function of r_+ for large r_+ in Fig.2a and Fig.2b.



Fig. 1

Figure 1. *Convergence in $Im(\omega)$ as order N is increased, for $r_+ = 10$*

It is seen that the both the imaginary and real parts of the mode frequency are proportional to r_+ for *large* r_+. The real and imaginary parts of the mode frequency can also be plotted as a function of the surface gravity α. It is seen that for *both* small and large r_+, the real and imaginary parts of the frequency are approximately proportional to α. $Re(\omega) \sim 0.24\ \alpha$ and $Im(\omega) \sim 0.41\ \alpha$ provides a good fit for the data. In the case of the QN modes where the modes could be obtained exactly, we saw that they were proportional to the surface gravity. Our numerical results seem to suggest that this may be true even for the five dimensional black hole - at least for very small and very large black holes. The surface gravity $\alpha \;=\; (2r_+ \;+\; \frac{1}{r_+})$. Therefore, α is large for very small and very large black holes. We have verified numerically that for very small black holes, the mode frequencies are very large, and $\omega \sim \frac{1}{r_+}$. This behaviour of the mode frequency is expected - as can be seen from the differential equation (27) on changing to the inverse radial coordinate x given by (29). Then, the differential equation in these coordinates is

$$(-r_0^2x^6 + x^4 + x^2)\frac{d^2\psi}{dx^2} \;+\; (2x^3 - 4r_0^2x^5 + 2i\omega x^2)\frac{d\psi}{dx} \;-\; (\frac{15}{4} + \frac{3x^2}{4} + \frac{9r_0^2x^4}{4})\psi \;=\; 0 \qquad (38)$$



Fig. 2a

Figure 2a. *$Re(\omega)$ as a function of r_+ for large r_+*

Now, we scale x as $x \;=\; q\ x_+$ (where $x_+ = \frac{1}{r_+}$). Now, (38) is

$$(-x_+^2 q^6 - q^6 + x_+^2 q^4 + q^2)\frac{d^2\psi}{dq^2} \;+ \\ (2x_+^2 q^3 - 4x_+^2 q^5 - 4q^5 + 2i\omega x_+ q^2)\frac{d\psi}{dq} \;- \\ (\frac{15}{4} + \frac{3x_+^2 q^2}{4} + \frac{9x_+^2 q^4}{4} + \frac{9q^4}{4})\psi \;=\; 0 \tag{39}$$

Then we see that for very large black holes (small x_+ approximation), x_+ can be scaled away from (39) near the horizon in this approximation provided $\omega \;=\; \frac{C}{x_+}$ where C is a constant independent of x_+. It can also be checked that near the horizon, there are no solutions to the scaled equation with $C \;=\; 0$, except the trivial solution. This shows that ω is indeed proportional to r_+ (i.e $\frac{1}{x_+}$) for large black holes.

For very small black holes (i.e in the large x_+ approximation), again x_+ can be scaled away



Fig. 2b

Figure 2b. *$Im(\omega)$ as a function of r_+ for large r_+*

from (39) near the horizon provided $\omega = D\ x_+$ where D is a constant independent of x_+. Here too, it can be checked that near the horizon, there are no solutions to (39) in the large x_+ approximation with $D = 0$ (except the trivial solution). This implies that for small black holes, ω is proportional to $\frac{1}{r_+}$ (i.e to x_+). This is also just a reflection of the fact that a very small AdS-Schwarzschild black hole has negative specific heat and actually resembles a Schwarzschild black hole.

We now examine the previous numerical work on QN modes of AdS black holes in four, five and seven dimensions.

The numerical results obtained in [7] for the five dimensional black hole do not agree with our values for the QN modes. There, the solution to (27) is expanded as a series around the horizon in the inverse radial coordinate $x = 1/r$. Using this, the l.h.s of (27) is expanded as a series around the horizon, and a recursion relation is obtained for the coefficients in the expansion of the solution. The coefficients a_n are functions of the black hole parameters and



Fig. 3a

Figure 3a. *Re(ω) as a function of α*

the mode frequency ω. Therefore,

$$\psi = \sum_n a_n(\omega)\ (x - x_+)^n \tag{40}$$

The mode frequencies are then the roots of the equation obtained by setting the series to zero at $x = 0$. In actual computation, the series is truncated, and the roots obtained. However, the mode frequencies seem to go to zero as $r_+ \to 0$, which is not the behaviour expected from the differential equation.



Fig. 3b

Figure 3b. *Im(ω) as a function of α*

A numerical computation of QN modes is tricky because of the nature of the boundary conditions on the mode solution at both the horizon and at infinity. The mode, which is ingoing (and not normalisable) at the horizon, could be contaminated by an outgoing component which goes to zero there. At the other boundary, the solution is required to go to zero. In a numerical truncation, it could also be contaminated as $x \to 0$ (i.e as $r \to \infty$) by the solution that blows up at this end. Therefore, in a numerical computation, it is essential to ensure the correct ingoing behaviour near the horizon and the correct *asymptotic* behaviour of the solution that goes to zero as $r \to \infty$. In our method, similar to the continued fraction method for asymptotically flat black holes, we have a *specific* ansatz for the wave function where the behaviour at *both* boundaries is explicitly present in the form of the ansatz. The form of our ansatz ensures that there is no contamination from the outgoing part at the horizon. At the other end as $x \to 0$ (i.e as $r \to \infty$), we *demand* that the solution must fall off as $x^{\frac{5}{2}}$ at every order. This *ensures* that there is no contamination from the solution that blows up at this end.

We have presented a new approach to computing the quasi-normal modes of *AdS* black holes. The novel feature of this method was an ansatz for the QN mode wave function which was derived from a superpotential. This was made in analogy with the case of the three dimensional BTZ black hole where the modes can be obtained exactly and the wave function is derived from a superpotential. The BTZ QN modes were proportional to the surface gravity. The modes obtained by us numerically for the five dimensional black hole are also approximately proportional to the surface gravity. More importantly, the modes obtained by us numerically are proportional to the inverse of the horizon radius for small black holes, reflecting the well-known fact that these black holes are unstable. We have also shown, from some scaling properties of the differential equation obeyed by the mode solutions, that this is indeed to be expected for small black holes. Work is in progress to compute the QN modes for AdS black holes in four and seven dimensions, and also the corresponding Reissner-Nordstrom black holes using this approach. Some preliminary work on the five dimensional RN AdS black hole seems to suggest that the QN mode increases with the charge of the black hole.

We would like to thank G.T.Horowitz and V.Hubeny for useful comments about the results in our paper.

References

[1] C.V.Vishveshvara, Nature **227**, 936 (1970)

[2] H.Nollert, Class.Quant.Grav. **16**, R159 (1999)

[3] K.D.Kokkotas, B.G.Schmidt, Living Reviews in Relativity, **2** (1999)

[4] E.Witten, Adv.Theor.Math.Phys. **2**, 253 (1998)

[5] U.H.Danielsson, E.Keski-Vakkuri and M.Kruczenski, JHEP **0002**, 039 (2000)

[6] S.Kalyana Rama and B.Sathiapalan, Mod.Phys.Lett **A14**, 2635 (1999)

[7] G.T.Horowitz and V.Hubeny, Phys.Rev.D **62**, 024027 (2000)

[8] T.R.Govindarajan, V.Suneeta and S.Vaidya, Nucl.Phys.B **583**, 291 (2000), and references therein.

[9] E.Witten, Adv.Theor.Math.Phys.**2**, 505 (1998)

[10] S.Chandrasekhar, *The Mathematical Theory of Black Holes*, Clarendon Press, Oxford (1983) and references therein.

[11] H-P. Nollert and B.G.Schmidt, Phys.Rev.D **45**, 2617 (1992)

[12] B.F.Schutz and C.M.Will, Astrophys. J. **291**, L33 (1985)

[13] E.W.Leaver, Proc.Roy.Soc.London, Ser.A **402**, 285 (1985)

[14] V.Ferrari and B.Mashhoon, Phys.Rev.D **30**, 295 (1984)

[15] H-J. Blome and B. Mashhoon, Phys. Lett. **A 100**, 231 (1983)

[16] M.Banados, C.Teitelboim and J.Zanelli, Phys.Rev.Lett. **69**, 1849 (1992)

[17] F.Cooper, A.Khare and U.Sukhatme, Phys.Rep **251**, 267 (1995)